

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/274511501>

A study on the geographical distribution along with habitat aspects of rodent species in Turkey

Article in Bonn Zoological Bulletin · January 2003

CITATIONS

13

READS

61

4 authors:



Nuri Yiğit

Ankara University

157 PUBLICATIONS 1,477 CITATIONS

[SEE PROFILE](#)



Ercüment Çolak

Ankara University

178 PUBLICATIONS 1,613 CITATIONS

[SEE PROFILE](#)



Mustafa Sözen

Bülent Ecevit Üniversitesi

158 PUBLICATIONS 1,287 CITATIONS

[SEE PROFILE](#)



Önder Özkurt

Ahi Evran Üniversitesi

63 PUBLICATIONS 601 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Ankara Üniversitesi Tandoğan Kampüsü Biyoçeşitliliğinin Belirlenmesi Projesi [View project](#)



Determining Genetic Variations and Phylogenetic Relationships of Genus Microtus and Chionomys Species (Mammalia: Rodentia) Distributed in Northern Turkey [View project](#)

A study on the geographical distribution along with habitat aspects of rodent species in Turkey

N. Yiğit, E. Çolak, M. Sözen & S. Özkurt

A b s t r a c t. Rodent species were collected in 20 selected localities, and these localities were then compared with regard to rodent species composition. A total of 41 rodent species was found in these localities, the number in each locality ranging from 9 to 17. Species previously recorded in mixed and deciduous forest in northern Asiatic Turkey, such as *Glis glis*, *Muscardinus avellanarius*, *Microtus subterraneus*, *Microtus roberti*, *Microtus majori*, *Sciurus vulgaris* and *Clethrionomys glareolus*, were not found in forested localities in west and south Asiatic Turkey with a dry summer season. *Apodemus agrarius* was only recorded from the Thracian region of Turkey. Similar habitats in different localities supported rodent assemblage with considerable differences in species composition. Vegetation structure, climate, and elevation were found to be the main factors affecting the distribution of rodent species in Turkey.

K e y w o r d s. Rodent species, ecological analysis, biogeography, geographical ecology, Turkey.

Introduction

Turkey comprises 775.000 km² in Asia and 4.450 km² in Europe (Thrace), 779.450 km² in total. Corbet (1978) listed 217 rodent species in the Palaearctic region, including Turkey. Demirsoy (1996) stated in his brief review, ‘The mammals of Turkey’, that 61 rodent species occur in Turkey. The following authors have contributed to the known Turkish rodent fauna by providing distributions and records of new species and subspecies: Danford & Alston (1877); Thomas (1897, 1903, 1906 & 1919); Barret-Hamilton (1900); Nehring (1903); Satunin (1908); Miller (1908); Blackler (1916); Aharoni (1932); Neuhäuser (1936); Ellerman (1951); Misonne (1957); Kahmann (1961); Spitzemberger & Steiner (1964); Osborn (1962 & 1965); Steiner & Vauk (1966); Lehmann (1966 & 1969), Lay (1967); Felten & Storch (1968); Felten et al. (1971 & 1973); Mursaloğlu (1965 & 1973); Spitzemberger (1971 & 1978); Morlok (1978); Kivanç (1983 & 1986); Doğramacı (1989); Doğramacı et al. (1994); Kurtonur & Özkan (1991); Çolak & Kivanç (1991); Çolak et al. (1994, 1997a, 1997b, 1998 & 1999); Kefelioğlu (1995); Filipucci et al. (1996); Coşkun (1996 & 1997); Yiğit et al. (1997a, 1997b, 1998a 1998b & 1998c); Yiğit & Çolak (1999); Özkurt et al. (1999). The number of rodent species in Turkey is now 61 according to the latest records. Although many rodent species have wide distribution areas in Turkey, information on geographical distributions, habitat peculiarities and ecology are insufficient. However, many studies on various aspects of rodent population structure, zoogeography, and ecology have been conducted in neighbouring areas by Thomas (1905); Zahavi & Wahrman (1957); Bodenheimer (1958); Misonne (1959); Hatt (1959); Ondrias (1966); Haim & Tchernov (1974); Atallah (1977);

Brown (1980). Land-use activities in Turkey such as farming in natural steppe areas; urban building and forestry dangerously affect the preservation of natural condition for wildlife. That is why rodent species, like other mammal species, are threatened with extinction. The aims of this study were to inform authorities establishing conservation areas, and to contribute to the composition of species and distribution areas of rodent species in Turkey, as well as further investigations.

Materials and Methods

This study, carried out between the years 1991 and 1997, is based on specimens caught in various localities in Turkey (Fig. 1). These were generally the type localities and record area of certain species, and detailed descriptions are provided here from 20 localities. When there were different habitat types in a selected locality, they were studied separately, each one consisting of close to the 500 1 ha plots. Meteorological records such as precipitation, mean temperature of the coldest month (m), mean temperature of the warmest month (M) and elevation of each locality were obtained from the nearest meteorological station. In addition vegetation structure, and the latitude and longitude of the localities were determined. Field studies were usually conducted in spring and summer months. We worked in 20 localities in periods of 3 days between 1991–1997. 200 snap traps and 50 Sherman live traps were set in the field daily in the afternoon, and checked early in the morning so that trapped specimen would not be damaged by insects. Some of these traps were also set on tree branches to catch arboreal rodent species. Traps were baited with a mixture of roasted peanuts and bread. *Allactaga* species were caught with small insect nets thrown from a slowly moving car at night, and special traps were used to catch fossorial species. Four external characteristic measurements (head and body, tail, hind foot and ear) and body weight were taken in the field. Cranial examinations were used along with these measurements to identify species. Additionally, the checklists of Corbet (1978) and Harrison & Bates (1991) were referred to for the identification of these species. The similarity coefficients were computed in order to compare rodent species composition in the localities by using NTSYS-pc computer programme according to Rohlf (1988). Skin and skulls of specimens were deposited at the University of Ankara, Faculty of Science, Department of Biology.




Fig. 1. Map showing localities of Turkish rodents recorded in this study. 1: Velika, Kırklareli 2: Bayındır, İzmir, 3: Ovacık, İzmir, 4: Demirci, Manisa, 5: Çardak, Denizli, 6: Çığlıkara, Antalya, 7: Kilbasan, Karaman, 8: Göçekhisik, Eskisehir, 9: Abant, Bolu, 10: Yenikonak, Bursa, 11: Tosya, Kastamonu, 12: Yıldızeli, Sivas, 13: Türkoğlu, Kahramanmaraş, 14: Kilis 10 km east, 15: Ceylanpınar, Sanliurfa, 16: Darende, Malatya, 17: Van, 10 km south, 18: Aralık, İğdır, 19: Sümela, Trabzon, 20: Kars and Ardahan.

Abbreviations in text: **1. VEL:** Velika, Kırklareli, **2. BAY:** Bayındır, İzmir, **3. OVA:** Ovacık, İzmir, **4. DEM:** Demirci, Manisa, **5. ÇAR:** Çardak, Denizli, **6. ÇIG:** Çıglıkara, Antalya, **7. KIL:** Kılbasan, Karaman, **8. GOK:** Gökçekışık, Eskişehir, **9. ABA:** Abant, Bolu, **10. YEN:** Yenikonak, Bursa, **11. TOS:** Tosya, Kastamonu, **12. YIL:** Yıldızeli, Sivas, **13. TUR:** Türköglu, Kahramanmaraş, **14. KILS:** Kilis, 10 km east, **15. CEY:** Ceylanpınar, Şanlıurfa, **16. DAR:** Darende, Malatya, **17. VAN:** Van, 10 km south, **18. ARA:** Aralık, İğdır, **19. SUM:** Sümela, Trabzon, **20. KA:** Kars and Ardahan.

Results and Discussion

During this study, 41 rodent species were captured in 20 localities in Turkey (Fig. 1). Of these, *Cricetulus migratorius*, *Rattus rattus*, and *Mus aff. musculus*, occurring in all localities studied, were identified as wide-ranging species. *Rattus norvegicus* were not trapped in localities studied that consist of rural areas, but this species is very common around urban areas (Yiğit et al. 1998a). *Apodemus hermonensis* and *Dryomys nitedula* were recorded in 16 and 15 different localities, respectively (Table 1). *A. hermonensis* was first reported in western Turkey by Filipucci et al. (1996). The following species, previously recorded from Turkey (Danford & Alston 1877; Thomas 1897; Thomas 1905; Neuhäuser 1936; Misonne 1957; Spitzemberger 1971 & 1978; Mursaloğlu 1973; Çolak & Kivanç 1991; Doğramacı et al. 1994; Kivanç et al. 1997a), were not taken in any of the localities studied: *Spermophilus citellus*, *Microtus arvalis*, *Microtus gud*, *Arvicola terrestris*, *Micromys minutus*, *Myomyscus roachi*, *Eliomys quercinus*, *Dryomys pictus*, *Meriones persicus*, *Meriones libycus*, *Tatera indica*, *Nesokia indica*, *Acomys cilicicus*, *Calomyscus bailwardi* and *Myocastor coypus*. Another species, *Spalax nehringi*, was recently recorded from Turkey by Coşkun (1996); its occurrence and taxonomic status are under discussion. We did capture *S. citellus*, *A. terrestris*, *M. persicus*, *A. cilicicus* and *Rattus norvegicus* in other localities of Turkey, but not in localities described in this article (Kivanç et al. 1997a; Yiğit & Çolak 1999; Özku et al. 1999; Yiğit et al. 1998a). In addition, we were told of the occurrence of *Castor fiber* in some virgin rivers in the coastal region of northern Anatolia by our colleagues and villagers.

Locality 11 had the highest rodent number with 17 species, followed by localities 9, 14 and 15 with 16 species each; locality 9 had the lowest rodent number with nine species (Table 1). Although we carried out thorough field studies in locality 14, we did not capture *Meriones sacramenti*, *M. libycus*, *M. vinogradovi*, *Nesokia indica* and *Tatera indica*, all recorded at locality 14 by Misonne (1957). Zahavi & Wahrman (1957) stated that *M. sacramenti* is endemic to Israel and was wrongly reported from this locality.

The similarity coefficient between the localities ranged from 1 to 0.512 (Table 2). Cluster analyses showed that the 20 localities studied formed three main clusters: the first is the northern localities (1, 9, 10, 19); the second is the western and central Anatolian localities (2, 3, 4, 5, 6, 7, 8, 11, 12, 16, 17, 18, 20); and the third is the south-eastern localities (13, 14, 15) (Fig. 2). The similarity coefficient ranged from 0.88 to 0.78 among localities of first main clusters, which are completely covered with mixed and deciduous forest. M and m values among these localities are also similar to each other (Tables 2, 3, 4). Seven rodent species (*Sciurus anomalus*, *C. migratorius*, *D. nitedula*, *Glis glis*, *Apodemus flavicollis*, *R. rattus*, *M. aff. musculus*) commonly occur in these four localities (Table 1). The main differences were the

Rodent species trapped and their distribution according to the localities (+ : occurrence status).

Table 1. ctd.

Rodent Species	13	12	11	14	12	15	16	13	17	12	9	16	15	14	13	15	14
<i>Microtus majori</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Promethomys schaposchnikowi</i>	-	+	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Ellobius latescens</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Spalax leucodon</i>	-	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-
<i>Spalax ehrenbergi</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Dryomys nitedula</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-
<i>Dryomys laniger</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Muscardinus avellanarius</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Glis glis</i>	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Apodemus sylvaticus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Apodemus flavicollis</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-
<i>Apodemus agrarius</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Apodemus mystacinus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-
<i>Apodemus hermannensis</i>	-	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-
<i>Apodemus uralensis</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Rattus rattus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-
<i>Mus aff. musculus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-
<i>Sicista caucasica</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Allactaga williamsi</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Allactaga euphratica</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Allactaga elater</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Hystrix indica</i>	-	+	+	-	+	-	-	-	-	-	-	-	-	-	-	-	-
Total numbers	13	12	11	14	12	15	16	13	17	12	9	16	15	14	13	15	14




Fig. 2. UPGMA clustering of Simqual matrix for rodent composition of 20 localities studied.

occurrence of *Microtus subterraneus* and *Apodemus agrarius* in locality 1 and *Microtus roberti* and *M. majori* in locality 19. A detailed report was provided for *M. subterraneus* and *M. majori* by Çolak et al. (1998), contributing to the distribution of *M. majori* and *M. subterraneus* in northern Anatolia, and it noted that *M. majori* does not occur in Abant (locality 9). The first and second main clusters therefore joined at similarity 0.68.

The second main clusters consist of localities selected in western, central and eastern Turkey (Fig. 1). The similarity coefficient among these localities ranged from 0.95 to 0.73 (Table 2). This group is composed of four sub-clusters. The localities 2, 3, 4 and 6 constitute the first sub-cluster and are located in the coastal region of western and south-western Asiatic Turkey, respectively. When localities 2, 3, 4 were compared to locality 6, the four localities joined at similarity 0.85 to form sub-clusters (Table 2). The climatic and vegetation peculiarities of these localities are given in tables 3, 4. Although the climatic data and vegetation were very similar among these localities, *Dryomys laniger* and *Microtus nivalis* were only recorded in locality 6, the former being an endemic rodent species for Asiatic Turkey. The factors restricting its distribution in other forest areas are still unknown. *M. nivalis*, *M. guentheri* and *D. nitedula* were previously reported in locality 6 by Spitzenerger (1978) and Kivanç et al (1997b). Steppe species such as *Cricetulus migratorius*, *Meriones tristrami* and *Allactaga williamsi* were caught in some parts of locality 4 where the forest had been cleared. Locality 2 is also at the western edge of the distribution of *M. tristrami* (Thomas 1905 & 1919). According to our finding, *Clethrionomys glareolus*, *Microtus subterraneus*, *M. majori*, *Glis glis* and *Muscardinus avellanarius*, which are distributed in the northern localities, do not occur in the localities 2, 3, 4 and 6 which have very dry summer months (Table 3). We suspect

that differences in climatic data, especially in precipitation, cause them to be absent from these localities.

The second sub-clusters consist of localities 5, 7, 8 in central and 11 in northern Anatolia (Fig. 1). The similarity coefficients vary between 0.95 and 0.89 in these localities (Table 2). Localities 5, 7 and 8 are characterized by steppe, and so the flora is composed mainly of steppe plants presented in table 4. However, some parts of locality 8 are covered with oak forest and bushes. Although northern Anatolia is generally covered with mixed forest, grain and rice fields are very abundant in locality 11. Because of this, steppe species such as *M. tristrami*, *A. williamsi* and *Mesocricetus brandti* manage to penetrate into this locality. Thus, locality 11 was determined to be near their range boundary in north-west Asiatic Turkey (Çolak et al. 1994; Yiğit et al. 1998b). In addition, we caught *S. anomalus* in locality 8 which has some forest areas. The climatic data of localities 5, 7 and 8 were very similar, but the *m* value of locality 5 was considerably higher than in localities 7 and 8. The main climatic difference between localities 5, 7, 8 and 11 is precipitation (Table 3). When these localities were compared with localities 12, 17, 16, which constitute the third sub-clusters and are also characterized by steppe (Fig. 1), the similarity coefficient between localities 5, 7, 8 and 11 and locality 12, 16 and 17 joined at similarity 0.85 (Table 2). The main differences among these localities are the occurrence of *Ellobius lutescens* in locality 17, and the absence of *M. tristrami* in localities 12, 17. *E. lutescens* was previously recorded from locality 17 by Coşkun (1997). The high mountains that extend diagonally from south Anatolia to north-east Anatolia should be considered as the main factors preventing *E. lutescens* from penetrating into central Anatolia. Although extensive field studies have been carried out in localities 12 and 17, we did not catch *M. tristrami* in these localities. The factor restricting *M. tristrami*'s distribution in these localities is still unclear.

Table 2. Simqual similarity matrix for qualitative data of 20 localities studied (see Table 3).

1.000																					
0.780	1.000																				
0.804	0.926	1.000																			
0.731	0.951	0.926	1.000																		
0.682	0.804	0.780	0.853	1.000																	
0.682	0.853	0.878	0.853	0.707	1.000																
0.658	0.829	0.804	0.878	0.926	0.780	1.000															
0.707	0.829	0.804	0.878	0.926	0.780	0.951	1.000														
0.780	0.756	0.780	0.756	0.756	0.707	0.682	0.731	1.000													
0.804	0.731	0.804	0.731	0.634	0.731	0.658	0.707	0.878	1.000												
0.707	0.780	0.756	0.829	0.926	0.682	0.853	0.902	0.829	0.707	1.000											
0.682	0.707	0.780	0.756	0.853	0.756	0.878	0.878	0.658	0.682	0.780	1.000										
0.658	0.829	0.804	0.829	0.780	0.780	0.756	0.707	0.682	0.658	0.707	0.682	1.000									
0.536	0.707	0.682	0.707	0.609	0.658	0.634	0.585	0.512	0.536	0.536	0.609	0.829	1.000								
0.536	0.707	0.682	0.707	0.609	0.658	0.634	0.585	0.512	0.536	0.536	0.609	0.829	1.000								
0.707	0.780	0.804	0.829	0.878	0.731	0.853	0.902	0.682	0.658	0.853	0.926	0.707	0.634	0.634	1.000						
0.682	0.707	0.780	0.756	0.853	0.707	0.829	0.829	0.658	0.634	0.780	0.951	0.682	0.609	0.609	0.926	1.000					
0.609	0.682	0.658	0.682	0.829	0.585	0.756	0.756	0.585	0.512	0.756	0.780	0.707	0.536	0.536	0.804	0.780	1.000				
0.804	0.731	0.804	0.731	0.682	0.682	0.658	0.707	0.829	0.853	0.756	0.682	0.658	0.536	0.536	0.707	0.682	0.560	1.000			
0.682	0.658	0.682	0.658	0.756	0.609	0.682	0.731	0.609	0.536	0.731	0.804	0.634	0.512	0.512	0.829	0.804	0.780	0.585	1.000		

Table 3. Average monthly precipitation by season and M and m values of the localities.

The localities and their numbers	Precipitation mm ³ /m ²					Climatic data M. m values
	Winter	Spring	Summer	Autumn	Total	
1. Velika / Kırklareli	102, 4	59,4	31,9	78,4	815,4	Rainy, temperate type Mediterranean climate. M: 34,4, m: 4.4
2. Bayındır / İzmir	125	50	6.3	47	684	Less rainy, temperate type Mediterranean climate. M: 34.4, m: 4.4
3. Ovacık / İzmir	129	43	1.5	45	655	Less rainy, cold type Mediterranean climate. M: 28, m: -2.2
4. Demirci / Manisa	109	64	13	49	705	Less rainy, cold type Mediterranean climate. M: 25, m: -3.1
5. Çardak / Denizli	51	44	15	24	402	Semiarid, cold type Mediterranean climate. M: 31,9, m: 0.2
6. Çığlıkara / Antalya	96	38	13	29	528	Semiarid, cold type Mediterranean climate. M: 31,9, m: 0.2 ***
7. Kılbasan / Karaman	36	34	10	20	300	Semiarid, very cold type Mediterranean climate. M: 30.3, m: -3.1
8. Gökçekisik / Eskişehir	43	41	19	24	381	Semiarid, very cold type Mediterranean climate. M: 28.4, m: -3.5
9. Abant / Bolu	57	53	34	38	546	Semiarid, very cold type Mediterranean climate. M: 25.7, m: -3.1
10. Yenikonak / Bursa	160	110	37	89	1184	Rainy,cold type Mediterranean climate. M: 25, m: -1.8
11. Tosya / Kastamonu	50	51	28	27	468	Semi continental, semiarid cold type climate. M: 27,1, m: -2.5
12. Yıldızeli / Sivas	42	55	13	22	399	Semiarid, very cold type Mediterranean climate. M: 27,8, m: -7.7
13. Türkoğlu / Kahramanmaraş	157	77	3	34	813	Less rainy, cold type Mediterranean climate. M: 35.0, m: 0.9
14. Kilis 10 km east	93	50	3	30	528	Semiarid, cold type Mediterranean climate. M: 35,0, m: -1,6
15. Ceylanpınar / Şanlıurfa	54	39	0.7	16	330	Semiarid, cold type Mediterranean climate. M: 40.6, m: 0.5
16. Darende / Malatya	42	52	11	43	383	Semiarid, very cold type Mediterranean climate. M: 29.9, m: -5.1
17. Van 10 km south	34	49	9	35	381	Semiarid,very cold type Mediterranean climate. M: 28.2, m:-8.0
18. Aralık / İğdır	15	28	10	13	198	Arid, cold type Mediterranean climate. M: 32.7, m: -8.4
19. Sümela / Trabzon	57.9	77.7	46.0	57.7	718.8	Less rainy, cold type Mediterranean climate. M: 25, m: -3.1
20. Kars and Ardahan	19.4	48.9	70.1	31	508.6	Less rainy, cold type Mediterranean climate. M: 25, m: -3.1

The fourth sub-cluster consists of localities 18, 20, and the similarity coefficient is 0.78 between localities 18 and 20, which are close to each other (Fig. 1; Table 2). This value is comparatively low, the main reason for the difference arose from locality 18. Locality 18 is interesting and its habitat is classified principally in two categories (Table 4): sandy plains toward the hillside, and marshy plains, sometimes completely flooded in the winter months. The vegetation structure of these habitat types is very different (Table 4). *Meriones meridianus*, *M. vinogradovi* and *Allactaga elater* are only present there. Indeed *M. meridianus* and *M. vinogradovi* were first recorded from Aralik by Yiğit et al. (1998c), and a subspecies of *A. elater* (*A. e. aralychensis*) was first described from Aralik by Satunin (1908). Another species of the genus *Allactaga* (*A. williamsi*) also occurs in this locality, ranging from the east to the west, except in localities 14 and 15 in southeastern Turkey. *M. meridianus* was caught only in sandy areas, and was found not to be penetrating into wet plains in locality 18. In contrast, *M. tristrami*, *M. vinogradovi* and *A. elater* were not caught in the sands, although *A. elater* is known to inhabit sandy areas (Shenbrot et al. 1994; Vinogradov & Argyropulo 1941). In addition, Vinogradov & Argyropulo (1941) and Heptner (1975) have described *M. meridianus* as inhabiting sandy areas, which is consistent with our findings. Detailed taxonomic studies were also performed on *M. meridianus*, *M. vinogradovi*, *A. elater* and *A. williamsi* by Yiğit et al. (1997b & 1998c); Çolak et al. (1994 & 1997a). In localities 18 and 20, *Mesocricetus brandti*, *Sicista caucasica*, *Sciurus anomalus*, *S. vulgaris*, *Microtus socialis* and *Prometheomys schaposchnikowi* are distributed only in locality 20, and not in locality 18. *P. schaposchnikovi* was recorded from two different localities in north-east Turkey by Çolak et al. (1999). In addition to this, Demirsoy (1996) noted, in his brief revision, that *S. caucasica* is distrubed in north-east Turkey, and his report is consistent with our findings. These differences in species composition were probably due to elevation and climate (Table 3, 4). Annual precipitation and elevation in these localities are very different from each other; locality 20 is covered with snow for long periods during the winter unlike locality 18. Second and third main clusters join at similarity 0.64.

The third main cluster is composed of the localities 13, 14, and 15 in south-eastern Turkey (Fig. 1). Localities 14 and 15 have a similarity coefficient of 1 with each other and of 0.83 with locality 13 (Table 2, Fig. 2). Locality 13 had the lowest rodent number with nine species (Table 2). Although localities 13, 14 and 15 form a cluster, and there is geographic proximity among these localities, the rodent composition and climatic data are very different between locality 13 and localities 14, 15 (Table 1, 3, 4). Localities 14 and 15 are surrounded by highland areas in the north with an elevation of over 1000 m. The average annual precipitation of locality 13 is very different from that of localities 14 and 15, which have a dry summer period (Table 3). Thus, species accustomed to semi-arid conditions were found in localities 14 and 15. *Meriones crassus*, *Gerbillus dasyurus*, *Microtus irani*, *Spalax ehrenbergi* and *Allactaga euphratica* were only recorded from these localities. Whereas species that range over the central Anatolia steppes, such as *A. williamsi*, *M. brandti*, *Spermophilus xanthoprymnus*, *Spalax leucodon* and *Microtus epiroticus* were not caught in localities 13, 14 and 15. *M. crassus* and *G. dasyurus* were first recorded from localities 14 and 15 by Yiğit et al. (1998c & 1997a). Additionally, the taxonomic status of *A. euphratica* and *M. irani* was revealed by Çolak et al. (1994 & 1997b). Our findings agree with the habitats specified by Haim & Tchernov (1974); Brown (1980)

Table 4. Some geographical and ecological remarks on localities in which the studies were performed. La.: Latitude, Lon.: longitude, El.: Elevation (m).

The localities	La.	Lon.	El.	The peculiarity of the localities and dominant plant species
1. Velika / Kırklareli	41.49	27.46	800	This locality consists of mixed forest with <i>Phillyrea latifolia</i> , <i>Fagus orientalis</i> , <i>Quercus pubescens</i> , <i>Quercus cerris</i> .
2. Bayındır / İzmir	38.13	37.39	100	This locality partly consists of cultivated areas, open hill sides, olive trees and shrubs (<i>Pinus brutia</i> , <i>Cistus creticus</i> , <i>Rhus coriaria</i> , <i>Olea europaea</i> var. <i>oleaster</i> , <i>Pistacia lentiscus</i> , <i>Quercus coccifera</i>)
3. Ovacık / İzmir	38.8	27.45	1150	This locality was dominantly covered with pines (<i>Pinus brutia</i> , <i>Cistus creticus</i> , <i>Rhus coriaria</i> , <i>Pistacia terebinthus</i>)
4. Demirci / Manisa	39.03	28.39	1350	This locality has the forested areas with pine, fruits and open fields in forest (<i>Pinus nigra</i> ssp. <i>pallasiana</i> , <i>Quercus cerris</i> , <i>Q. pubescens</i>)
5. Çardak / Denizli	37.50	29.40	920	This locality studied consists of grain fields and steppe (<i>Astragalus</i> sp., <i>Medicago radiata</i> , <i>Festuca</i> sp., <i>Cynodon</i> sp., <i>Thymus</i> sp., <i>Polygonum</i> sp.)
6. Çıglıkara / Antalya	36.43	29.55	1250	This locality was partly covered with mixed forests (<i>Pinus nigra</i> ssp. <i>pallasiana</i> , <i>Cedrus libani</i> , <i>Juniperus excelsa</i> , <i>Juniperus oxycedrus</i>)
7. Kılbasan / Karaman	37.19	33.11	1050	This locality consists of grain fields and steppe (plant species is the same as in Cardak / Denizli)
8. Göçekisksik / Eskişehir	39.39	30.20	900	This locality consists of grain fields and steppe (<i>Salvia cryptantha</i> , <i>Thymus spyleus</i> , <i>Ziziphora capitata</i> , <i>Teucrium polium</i>)
9. Abant / Bolu	40.40	31.45	1100	This locality is utterly covered with mixed forest (<i>Abies nordmanniana</i> , <i>Fagus orientalis</i> , <i>Corimus betulus</i> , <i>Populus tremula</i> , <i>Quercus infectoria</i> , <i>Q. cerris</i>)
10. Yenikonak / Bursa	40.07	29.10	1025	This locality is utterly covered with mixed forest (vegetation structure is the same as in Abant / Bolu)
11. Tosya / Kastamonu	41.01	34.02	870	This locality consists of mixed forest, shrubs and grain fields (<i>Quercus pubescens</i> , <i>Cistus laurifolius</i> , <i>Crataegus monogyna</i> , <i>Cotonaastere nummalaria</i>).
12. Yıldızeli / Sivas	39.52	36.36	1415	This locality is high steppe with <i>Astragalus angustifolius</i> , <i>Salvia aethiopis</i> , <i>Senecio vernalis</i> , <i>Hyoscyamus niger</i> .
13. Türkoğlu / Kahramanmaraş	37.24	36.61	520	This locality is composed of wetland, corn fields and swamp (<i>Cyperus longus</i> , <i>Carex otrubae</i> , <i>Bolbochoenus maritimus</i> , <i>Scilla bifolia</i>).
14. Kilis 10 km east	36.43	37.07	650	This locality has grain fields, natural steppe and rocky hills. Plant species is the same as in Ceylanpınar/Sanhurfa
15. Ceylanpınar / Şanlıurfa	36.51	40.03	400	This locality consists of cultivated areas and natural steppes (<i>Hordeum</i> sp., <i>Eryngium</i> sp., <i>Securigera</i> sp., <i>Peganum</i> sp., <i>Agropyron</i> sp.).
16. Darende / Malatya	38.33	37.31	1200	This locality has steppe and grain fields in roughness areas (<i>Centranthus longiflorus</i> , <i>Parietaria judaica</i> , <i>Torilis leptophylla</i> , <i>Eryngium campestre</i>).
17. Van 10 km south	38.27	43.19	1700	This locality includes high steppe and grain fields (<i>Festuca valesiaca</i> , <i>Eremopyea songarica</i> , <i>Bromus danthoniae</i> , <i>Ornithogalum</i> sp.).
18. Aralık / İğdır	39.53	44.31	825	This place is briefly composed of two types of localities. First is sandy, consists of following plant species: <i>Equisetum ramosissimum</i> , <i>Atraphaxis billardieri</i> , <i>Crucifera</i> sp., <i>Crepis</i> sp., <i>Medicago</i> sp., <i>Euphorbia</i> sp. Latter is watery, plains consist of following plant species: <i>Juncus</i> sp., <i>Dactylis</i> sp., <i>Alyssum</i> sp., <i>Erysimum</i> sp., <i>Carex</i> sp.
19. Sümela / Trabzon	40.47	39.37	1100	This locality is utterly covered with mixed forest (<i>Castanea sativa</i> , <i>Alnus glutinosa</i> , <i>Picea orientalis</i> , <i>Fagus orientalis</i> , <i>Juglans regia</i> , <i>Carpinus orientalis</i>).
20. Kars and Ardahan	41.07	42.43	1829	This locality is high steppe, and is dominantly covered with <i>Bromus tomentellus</i> , <i>Festuca valesiaca</i> , <i>Astragalus microcephalus</i> , <i>Agropyron repens</i> , <i>Echinops ritro</i> , <i>Eryngium campestre</i> .

and Shenbroth et al. (1994). We conclude that the cold climate and elevation affect the distribution of the species recorded from localities 14 and 15 into central Anatolia.

The similarity in the species compositions was found to vary greatly in localities with similar habitats such as forests, steppes, sands and wetlands. The temperature, elevation, precipitation and humidity were also found to vary from locality to locality, all of which are factors contributing to species composition in Turkey. The climatic diversity, geographic barriers that extend from south-eastern into northern Anatolia, and the connection of Anatolia with three continents, Europe, Asia and Africa, all resulted in faunistic and floristic diversity. Arboreal rodent species from the European continent, steppe species from Caucasia and arid-semi arid land species from Africa occupy different habitats in Anatolia. Niethammer & Krapp (1978 & 1982) stated that 61 rodent species live in the European continent. Considering Niethammer & Krapp (1978 & 1982), it can be said that Turkey is very rich in rodent species compared to the European continent.

Acknowledgements

This study was funded in part by the Research Fund of the Ankara University (No: 91250055, 91250072, 95050301, 96050306, 97050304, 98050308) and The Scientific and Technical Research Council of Turkey (TÜBÝTAK) (TBAG: 1186 and 1187). We thank Dr. Latif Kurt for identifying plant species and for his many helpful comments.

Zusammenfassung

An 20 ausgesuchten Orten der Türkei wurden Nagerarten gesammelt. Diese Orte wurden dann auf Artenzusammensetzung untersucht und miteinander verglichen. Insgesamt wurden 41 Nagerarten gefunden, 9 bis 17 an jedem Ort. Arten, die früher in Misch- und Laubwald in der nordasiatischen Türkei festgestellt worden waren, wie *Glis glis*, *Muscardinus avellanarius*, *Microtus subterraneus*, *Microtus roberti*, *Microtus majori*, *Sciurus vulgaris* und *Clethrionomys glareolus*, wurden in bewaldeten Gegenden der west- und südasiatischen Türkei mit trockenem Sommer nicht gefunden. *Apodemus agrarius* wurde nur im türkischen Thrakien verzeichnet. Ähnliche Habitate an verschiedenen Orten hatten Nageransammlungen mit beträchtlichen Unterschieden in der Artenzusammensetzung. Vegetation, Klima und Höhe wurden als Hauptfaktoren für die Verbreitung von Nagerarten in der Türkei ermittelt.

References

- Aharoni, B. (1932): Die Muriden von Palastina und Syrien. – Z. Säugetierk. 11: 161–240.
Atallah, S. I. (1977): Mammals of the eastern Mediterranean region: their ecology, systematics and zoogeographical relationships. – Säugetierk. Mitt. 26: 1–50.
Barrett-Hamilton, G. E. H. (1900): On geographical and individual variation in *Mus sylvaticus* and its allies. – Proc. Zool. Soc. Lond. 1900: 378–428, pl. 25.
Blackler, W. F. G. (1916): On a new species of *Microtus* from Asia Minor. – Ann. Mag. nat. Hist. 17: 426–427.
Bodenheimer, F. S. (1958): The present taxonomic status of the terrestrial mammals of Palestine. – Bull. Res. Counc. of Israel 7b: 165–190.
Brown, R. E. (1980): Rodents of the Kavir National Park, Iran. – Mammalia 44: 89–96.
Corbet, G. B. (1978): The mammals of the Palaearctic region: a taxonomic review. – Brit. Mus. Nat. Hist. & Cornell Univ. Press, London.
Coşkun, Y. (1996): A new subspecies of *Spalax nehringi* (Satunin, 1898) (Rodentia: Spalacidae) from Turkey. – Säugetierk. Mitt. 37: 103–109.
Coşkun, Y. (1997): Türkiye *Ellobius lutescens* Thomas, 1897 (Rodentia: Cricetidae) Türünün Morfolojik ve Karyolojik Özellikleri. – Tr. J. Zoology 21: 349–354.

- Çolak, E. & E. Kivanç (1991): Distribution and taxonomic status of genus *Clethrionomy* Tilesius, 1850 (Mammalia: Rodentia) in north Anatolia. – Commun. Fac. Sci. Univ. Serie C, V. 9: 1–16.
- Çolak, E., E. Kivanç & N. Yiğit (1994): A study on taxonomic status of *Allactaga euphratica* Thomas, 1881 and *Allactaga williamsi* Thomas, 1897 (Rodentia: Dipodidae) in Turkey. – Mammalia 58: 591–600.
- Çolak, E., E. Kivanç & N. Yiğit (1997a): *Allactaga elater aralychensis* (Satunin, 1901) in Taksonomik Durumu ve Yayılışı. – Tr. J. Zoology 21: 355–360.
- Çolak, E., N. Yiğit, M. Sözen & P. Özkurt (1997b): Distribution and taxonomic status of the genus *Microtus* (Mammalia: Rodentia) in south-eastern Turkey. – Israel J. Zoology, 43: 391–396.
- Çolak, E., N. Yiğit, M. Sözen & P. Özkurt (1998): A study on taxonomic status of *Microtus subterraneus* (de Selys Longchamps, 1836) and *Microtus majori* Thomas, 1901 (Mammalia: Rodentia), in Turkey. – Tr. J. Zoology 22: 119–129.
- Çolak, E., N. Yiğit & R. Verimli (1999): On the karyotype of the long-clawed mole rat *Prometheomys schapochnikovi* Satunin, 1901 (Mammalia: Rodentia) in Turkey. – Z. Säugetierk. 64: 1–2.
- Danford, C. G. & E. R. Alston (1877): On the mammals of Asia Minor. – Proc. Zool. Soc. London 270–282.
- Demirsoy, A. (1996): Türkiye Omurgalıları “Memeliler”. – Çevre Bakanlığı Çevre Koruma Genel Müdürlüğü Proje no: 90 – K – 1000 – 90, Ankara.
- Doğramacı, S. (1989): Türkiye Memeli Faunası için Yeni Bir Tür, *Microtus epiroticus* (Mammalia: Rodentia). – Tr. J. Zoology 13: 197–203.
- Doğramacı, S., H. Kefelioğlu & I. Gündüz (1994): Türkiye *Spermophilus* (Mammalia: Rodentia) Cinsinin Karyolojik Analizi. – Tr. J. Zoology 18 (3): 167–170.
- Ellerman, J. R. (1948): Key to the Rodents of south-west Asia in the British museum collection. – Proc. Zool. Soc. 118: 765–816.
- Felten, H. & G. Storch (1968): Eine neue Schläferart *Dryomys laniger* n. sp. aus Kleinasien (Rodentia: Gliridae). – Senckenberg. Biol. 49: 429–435.
- Felten, H., F. Spitznerberger & G. Storch (1971): Zur Kleinsäugerfauna West-Anatoliens, Teil 1. – Senckenberg. Biol. 52: 393–424.
- Felten, H., F. Spitznerberger & G. Storch (1973): Zur Kleinsäugerfauna West-Anatoliens, Teil 2. – Senckenberg. Biol. 54: 227–290.
- Filipucci, M. G., G. Storch & M. Macholan (1996): Taxonomy of the genus *Sylvaemus* in Western Anatolia – Morphological and electrophoretic evidence. – Senckenberg Biol. 75: 1–14.
- Haim, A. & E. Tchernov (1974): The distribution of myomorph rodents in the Sinai Peninsula. – Mammalia 38: 201–223.
- Harrison, D. L. & P. J. J. Bates (1991): The Mammals of Arabia. – Second Edition. Harrison Zool. Museum Pub. Kent, England.
- Hatt, R. T. (1959): The mammals of Iraq. – Miscellaneous Publs Mus. Zool. Univ. Mich. 106: 1–113.
- Heptner, W. G. (1975): Über einige Besonderheiten der Formbildung und der geographischen Verbreitung der Rennmaus, *Meriones (Pallasiomys) meridianus* Pallas, 1773. – Z. Säugetierk. 40: 261–269.
- Kahmann, H. (1961): Beiträge zur Säugetierkunde der Türkei. 2. Die Brandmaus (*Apodemus agrarius* Pallas, 1774) in Thrakien und die südeuropäische Verbreitung der Art. – Rev. Fac. Sci. Univ. Istanbul 26: 87–106.
- Kefelioğlu, H. (1995): Türkiye *Microtus* (Mammalia: Rodentia) Cinsinin Taksonomisi ve Yayılışı. – Tr. J. Zoology 19: 35–63.
- Kivanç, E. (1983): Die Haselmäuse, *Muscardinus avellanarius* L., in der Türkei. – Bonn zool. Beitr. 34: 419–428.
- Kivanç, E. (1986): *Microtus (Pitymys) majori* Thomas, 1906 in der europäischen Türkei. – Bonn. zool. Beitr. 37: 39–42.

- Kıvanç, E., M. Sözen, E. Çolak & N. Yiğit (1997a): Karyologic and phallic aspects of spiny mouse, *Acomys cilicicus* Spitzenberger, 1978 (Rodentia: Muridae) in Turkey. – Tr. J. Zoology 21: 167–169.
- Kıvanç, E., M. Sözen, E. Çolak & Yiğit, N. (1997b): Karyologic and phallic characteristics of *Dryomys laniger* Felten and Storch, 1968 (Rodentia: Gliridae) in Turkey. – Israel J. Zool. 43: 401–403.
- Kurtonur, C. & B. Özkan (1991): New records of *Myomimus roachi* (Bate, 1937) from Turkish Tharace (Mammalia: Rodentia: Gliridae). – Senckenberg. biol. 71: 239–244.
- Lay, D. M. (1967): A study of the mammals of Iran, resulting from the street Expedition of 1962–63. – Fieldiana Zool. 54: 1–282.
- Lehmann, E. von (1966): Taxonomische Bemerkungen zur Säugetierausbeute der Kumerloeveschens Orientreisen. – Zool. Beitr. 12: 251–317.
- Lehmann, E. von (1969): Eine neue Säugetieraufsammlung aus der Türkei im Museum Koenig (Kumerloewe-Reise 1968). – Zool. Beitr. 15: 299–327.
- Miller, G. S. (1908): New mammals from Asia Minor. – Ann. Mag. nat. Hist. 1: 102–103.
- Misonne, X. (1957): Mammifères de la Turquie Sub-orientale et du nord de la Syrie. – Mammalia 21: 53–57.
- Misonne, X. (1959): Analyse zoogéographique des mammifères de l'Iran. – Mem. Inst. Sci. Natur. Belg. Bruxelles 2: 1–157.
- Morlok, W. F. (1978): Nagetiere aus der Türkei (Mammalia: Rodentia). – Senckenberg. Biol. (3–4) 59: 155–162.
- Mursaloğlu, B. (1965): Geographic variation in *Citellus citellus* (Mammalia: Rodentia) in Turkey. – Türk Biol. Derg. 10: 78–109.
- Mursaloğlu, B. (1973): New records for Turkish rodents (Mammalia). – Communications (C) 17: 214–219.
- Nehring, A. (1903): Über eine Springmaus aus Nordwest-Kleinasien (*Allactaga williamsi laticeps*, n. subsp.). – Sitz. Ges. naturf. Freunde 4: 357–360.
- Neuhäuser, G. (1936): Die Muriden von Kleinasien. – Z. Säugetierk. 2: 161–236.
- Niethammer, J. & F. Krapp (1978): Handbuch der Säugetiere Europas. – Akademische Verlagsgesellschaft, Band I Rodentia I. Wiesbaden, Germany.
- Niethammer, J. & F. Krapp (1982): Handbuch der Säugetiere Europas. – Akademische Verlagsgesellschaft, Band 2/1 Rodentia II. Wiesbaden, Germany.
- Ondrias, J. C. (1966): The taxonomy and geographical distribution of the rodents of Greece. – Säugetierk. Mitt. 14: 1–136.
- Osborn, D. J. (1962): Rodents of subfamily Microtinae from Turkey. – J. Mammalogy 43: 515–529.
- Osborn, D. J. (1965): Rodents of subfamilies Murinae, Gerbillinae, and Cricetinae from Turkey. – The journal of the Egyptian Public Health Association XL: 401–421.
- Özkurt, P., E. Çolak, N. Yiğit, M. Sözen & R. Verimli (1999): Contribution to karyology and morphology of *Arvicola terrestris* (Lin., 1758) in Central Anatolia. – Tr. J. Zoology 23: 253–257.
- Rölf, J. F. (1988): NTSYS-pc Numerical Taxonomy and Multivariate Analysis System, Version 1.50. – Exeter Publishing LTD, New York.
- Satunin, K. A. (1908): Beiträge zur Kenntnis der Säugetierfauna Kleinasiens u. Transkaspiens. – Mitt. Kaukas. Mus. Tiflis 4: 42–141.
- Shenbrot, G. I., K. A. Rogovin & E. J. J. Heske (1994): Comparison of Niche-packing and Community Organisation in Desert Rodent in Asia and North America. – Aust. J. Zool. 42: 479–499.
- Spitzenberger, F. & H. Steiner (1964): *Prometheomys schaposchnikovi* in Nordost-Kleinasien. – Z. Säugetierk. 79: 116–124.
- Spitzenberger, F. (1971): Zur Systematik und Tiergeographie von *Microtus* (*Chinomys*) *nivalis* und *Microtus* (*Chinomys*) *gud* (Microtinae: Mammalia) in S. Anatolien. – Z. Säugetierk. 36: 370–380.
- Spitzenberger, F. (1978): Die Stachelmaus von Kleinasien, *Acomys cilicicus* n. sp. – Annals naturhist. Mus. Wien 81: 443–446.

- Steiner, H. & G. Vauk (1966): Säugetiere aus dem Beysehir Gebiet (Vilayet Konya Kleinasien). – Zool. Anz. 176: 97–102.
- Thomas, O. (1897): On two new rodents from Van, Kurdistan. – Ann. Mag. nat. Hist. (6) 20: 308–310.
- Thomas, O. (1903): On two new Muridae from Smyrna. – Ann. Mag. nat. Hist. (12) 7: 188–190.
- Thomas, O. (1905): On a collection of mammals from Persia and Armenia presented to the British Museum by Col. A. C. Bailward. – Proc. Zool. Soc. London 2: 519–527.
- Thomas, O. (1906): New Insectivores and Voles collected by Mr. A. Robert near Trabezond. – Ann. Mag. nat. Hist. 17: 418–419.
- Thomas, O. (1919): Notes on gerbils referred to the genus *Meriones*, with descriptions of new species and subspecies. – Ann. Mag. nat. Hist. 9: 263–273.
- Vinogradov, B. S. & A. I. Argyropulo (1941): Fauna of the USSR. Mammals. Key to the rodents (Translated from Russian). – Leningrad Pub., Moscow.
- Yiğit, N., E. Çolak, E. Kivanç, & M. Sözen (1997a): A new gerbil from Turkey: *Gerbillus dasyurus* Wagner, 1836 (Rodentia: Gerbillinae). – Israel J. Zool. 43: 13–18.
- Yiğit, N., E. Kivanç & E. Çolak (1997b): Türkiye'deki *Meriones* Illiger, 1811 (Mammalia: Rodentia) Türlerinin Teshis Karakterleri ve Yayılışı. – Tr. J. Zoology 21: 361–374.
- Yiğit, N., E. Çolak, M. Sözen & S. Özkurt (1998a): The taxonomy and karyology of *Rattus norvegicus* (Berkenhout, 1769) and *Rattus rattus* (Linnaeus, 1758), in Turkey. – Tr. J. Zoology 22: 203–212.
- Yiğit, N., E. Kivanç & E. Çolak (1998b): On the taxonomic status of *Meriones tristrami* Thomas, 1892 (Rodentia: Gerbillinae) in Turkey. – Zoology in the Middle East 16: 19–30.
- Yiğit, N., E. Kivanç & E. Çolak (1998c): Contribution to taxonomy and karyology of *Meriones meridianus* (Pallas, 1773) and *Meriones crassus* Sundevall, 1842 (Rodentia: Gerbillinae) from Turkey. – Z. Säugetierk. 63: 311–314.
- Yiğit, N. & E. Çolak (1999): A study of the taxonomy and karyology of *Meriones persicus* (Blanford, 1875) (Mammalia: Rodentia) in Turkey. – Tr. J. Zoology 23: 269–274.
- Zahavi, A. & J. Wahrman (1957): The cytotaxonomy, ecology and evolution of the gerbils and jirds of Israel (Rodentia: Gerbillinae). – Mammalia 2: 341–380.

Dr. Nuri Yiğit, Dr. Ercüment Çolak & Dr. Mustafa Sözen, University of Ankara Faculty of Science, Department of Biology, 06100 Besevler, Ankara, Turkey; Dr. Sakir Özkurt, Gazi University, Education Faculty of Kırşehir, Department of Biology, Kırşehir, Turkey.